

## Reaction Seminars 2021 (April 1, 2021)



**From one to many : a neutral history**

F. Miguel Marqués



*NATURE*

FEBRUARY 27, 1932

## Possible Existence of a Neutron

It has been shown by Bothe and others that beryllium when bombarded by  $\alpha$ -particles of polonium emits a radiation of great penetrating power, which has an absorption coefficient in lead of about  $0.3 \text{ (cm.)}^{-1}$ . Recently Mme. Curie-Joliot and M. Joliot found that the transference of energy to the proton was by a process similar to the Compton effect, and estimated that the beryllium radiation had a quantum energy of  $50 \times 10^6$  electron volts.

I have shown that the radiation ejects particles from hydrogen, helium, lithium, beryllium, carbon, air, and argon. The particles ejected from hydrogen behave, as regards range and ionising power, like protons with speeds up to about  $3.2 \times 10^9$  cm. per sec. The particles from the other elements have a large ionising power, and appear to be in each case recoil atoms of the elements.

These results, and others I have obtained in the course of the work, are very difficult to explain on the assumption that the radiation from beryllium is a quantum radiation, if energy and momentum are to be conserved in the collisions. The difficulties disappear, however, if it be assumed that the radiation consists of particles of mass 1 and charge 0, or neutrons.

The collisions of this neutron with the atoms through which it passes give rise to the recoil atoms, and the observed energies of the recoil atoms are in fair agreement with this view.

J. CHADWICK.

Cavendish Laboratory,  
Cambridge, Feb. 17.



*"But there was no doubt whatever in my mind, or I should not have written the Letter"*

*NATURE*

FEBRUARY 27, 1932

## Possible Existence of a Neutron

It has been shown by Bothe and others that beryllium when bombarded by  $\alpha$ -particles of polonium emits a radiation of great penetrating power, which has an absorption coefficient in lead of about  $0.3 \text{ (cm.)}^{-1}$ . Recently Mme. Curie-Joliot and M. Joliot found that the transference of energy to the proton was by a process similar to the Compton effect, and estimated that the beryllium radiation had a quantum energy of  $50 \times 10^6$  electron volts.

I have shown that the radiation ejects particles from hydrogen, helium, lithium, beryllium, carbon, air, and argon. The particles ejected from hydrogen behave, as regards range and ionising power, like protons with speeds up to about  $3.2 \times 10^9$  cm. per sec. The particles from the other elements have a large ionising power, and appear to be in each case recoil atoms of the elements.

These results, and others I have obtained in the course of the work, are very difficult to explain on the assumption that the radiation from beryllium is a quantum radiation, if energy and momentum are to be conserved in the collisions. The difficulties disappear, however, if it be assumed that the radiation consists of particles of mass 1 and charge 0, or neutrons.

The collisions of this neutron with the atoms through which it passes give rise to the recoil atoms, and the observed energies of the recoil atoms are in fair agreement with this view.

J. CHADWICK.

Cavendish Laboratory,  
Cambridge, Feb. 17.



*"But there was no doubt whatever in my mind, or I should not have written the Letter"*

NATURE

FEBRUARY 27, 1932

## Possible Existence of a Neutron

It has been shown by Bothe and others that beryllium when bombarded by  $\alpha$ -particles of polonium emits a radiation of great penetrating power, which has an absorption coefficient in lead of about  $0.3 \text{ (cm.)}^{-1}$ . Recently Mme. Curie-Joliot and M. Joliot found that the transference of energy to the proton was by a process similar to the Compton effect, and estimated that the beryllium radiation had a quantum energy of  $50 \times 10^6$  electron volts.

I have shown that the radiation ejects particles from hydrogen, helium, lithium, beryllium, carbon, air, and argon. The particles ejected from hydrogen behave, as regards range and ionising power, like protons with speeds up to about  $3.2 \times 10^9$  cm. per sec. The particles from the other elements have a large ionising power, and appear to be in each case recoil atoms of the elements.

These results, and others I have obtained in the course of the work, are very difficult to explain on the assumption that the radiation from beryllium is a quantum radiation, if energy and momentum are to be conserved in the collisions. The difficulties disappear, however, if it be assumed that the radiation consists of particles of mass 1 and charge 0, or neutrons.

The collisions of this neutron with the atoms through which it passes give rise to the recoil atoms, and the observed energies of the recoil atoms are in fair agreement with this view.

J. CHADWICK.

Cavendish Laboratory,  
Cambridge, Feb. 17.




*"But there was no doubt whatever in my mind, or I should not have written the Letter"*

- by "atoms" he means "nuclei" :

$$\rightarrow \epsilon_n \sim \text{few \% } (\bullet)$$

$$\rightarrow \epsilon_{xn} \approx (\epsilon_{1n})^x (\bullet\bullet\cdots\bullet)$$



NATURE

FEBRUARY 27, 1932

## Possible Existence of a Neutron

It has been shown by Bothe and others that beryllium when bombarded by  $\alpha$ -particles of polonium emits a radiation of great penetrating power, which has an absorption coefficient in lead of about  $0.3 \text{ (cm.)}^{-1}$ . Recently Mme. Curie-Joliot and M. Joliot found that the transference of energy to the proton was by a process similar to the Compton effect, and estimated that the beryllium radiation had a quantum energy of  $50 \times 10^6$  electron volts.

I have shown that the radiation ejects particles from hydrogen, helium, lithium, beryllium, carbon, air, and argon. The particles ejected from hydrogen behave, as regards range and ionising power, like protons with speeds up to about  $3.2 \times 10^9$  cm. per sec. The particles from the other elements have a large ionising power, and appear to be in each case recoil atoms of the elements.

These results, and others I have obtained in the course of the work, are very difficult to explain on the assumption that the radiation from beryllium is a quantum radiation, if energy and momentum are to be conserved in the collisions. The difficulties disappear, however, if it be assumed that the radiation consists of particles of mass 1 and charge 0, or neutrons.

The collisions of this neutron with the atoms through which it passes give rise to the recoil atoms, and the observed energies of the recoil atoms are in fair agreement with this view.

J. CHADWICK.

Cavendish Laboratory,  
Cambridge, Feb. 17.




*"But there was no doubt whatever in my mind, or I should not have written the Letter"*

- by "atoms" he means "nuclei" :


$$\rightarrow \epsilon_n \sim \text{few \% } (\bullet)$$

$$\rightarrow \epsilon_{xn} \approx (\epsilon_{1n})^x (\bullet\bullet\cdots\bullet)$$




$$\rightarrow \epsilon_{xn} < (\epsilon_{1n})^x \text{ due to "cross-talk" ...}$$

# The nucleus : a 'liquid drop'



$$B(N, Z) = N M_n + Z M_p - M(N, Z)$$




$$B(N, Z) = N M_n + Z M_p - M(N, Z)$$

► Semi-empirical liquid-drop formula :

15.67	$a_v A$	volume
17.23	$- a_s A^{2/3}$	surface
0.714	$- a_c Z^2 / A^{1/3}$	Coulomb
23.29	$- a_a (N - Z)^2 / A$	asymmetry
11.2	$\pm \delta / A^{1/2}$	pairing
[MeV]		

- multineutrons :  $Z = 0$ ?



$$B(N, Z) = N M_n + Z M_p - M(N, Z)$$

► Semi-empirical liquid-drop formula :


15.67	$a_v A$	volume
17.23	$- a_s A^{2/3}$	surface
0.714	$- a_c Z^2 / A^{1/3}$	Coulomb
23.29	$- a_a (N - Z)^2 / A$	asymmetry
11.2	$\pm \delta / A^{1/2}$	pairing
[MeV]		

• multineutrons :  $Z = 0$ ?

→ tetraneutron :  $B/A(4,0) = -17 \text{ MeV}!!!$


→ trineutron :  $B/A(3,0) = -22 \text{ MeV}!!!$





► Well-established facts:


- $N = 2$  (✗) ...  $10^{57}$  (✓)




► Well-established facts:

- $N = 2$  (X) ...  $10^{57}$  (✓)
- $({}^3\text{He})_2$  (X) ...  $({}^3\text{He})_N$  (✓):  $N \sim 30$

Guardiola, PRL 84 (2000) 1144







► Well-established facts:

- $N=2$  (✗) ...  $10^{57}$  (✓)
- $(^3\text{He})_2$  (✗) ...  $(^3\text{He})_N$  (✓):  $N \sim 30$


Guardiola, PRL 84 (2000) 1144



► Known  $M(N, Z)$ :




- $B(^5\text{H}) > 0!$  [ $M(4, 1) < 4m_n + m_p$ ]
- **LD**( $N \neq Z$ )?




► Well-established facts:

- $N=2$  (X) ...  $10^{57}$  (✓)
- $(^3\text{He})_2$  (X) ...  $(^3\text{He})_N$  (✓):  $N \sim 30$


Guardiola, PRL 84 (2000) 1144



► Known  $M(N, Z)$ :




- $B(^5\text{H}) > 0!$  [ $M(4, 1) < 4m_n + m_p$ ]
- **LD** ( $N \neq Z$ )? **LD** with surface-corr.  $\alpha_a \dots$
- “multineutron anomaly”?




► Well-established facts:

- $N=2$  (X) ...  $10^{57}$  (✓)
- $({}^3\text{He})_2$  (X) ...  $({}^3\text{He})_N$  (✓):  $N \sim 30$


Guardiola, PRL 84 (2000) 1144



► Known  $M(N, Z)$ :




- $B({}^5\text{H}) > 0!$  [ $M(4, 1) < 4m_n + m_p$ ]
- **LD** ( $N \neq Z$ )? **LD** with surface-corr.  $a_a \dots$
- “multineutron anomaly”?
- **even** neutron numbers:  $\boxed{{}_0^4\text{n}}$




► Well-established facts:

- $N=2$  (X) ...  $10^{57}$  (✓)
- $({}^3\text{He})_2$  (X) ...  $({}^3\text{He})_N$  (✓):  $N \sim 30$

Guardiola, PRL 84 (2000) 1144



► Known  $M(N, Z)$ :




- $B({}^5\text{H}) > 0!$  [ $M(4, 1) < 4m_n + m_p$ ]
- **LD** ( $N \neq Z$ )? **LD** with surface-corr.  $a_a \dots$
- “multineutron anomaly”?
- **even** neutron numbers:  ${}^4_0\text{n}$


► Two important issues:

- **production** (● unstable)
- **detection** (extremely low ●●●●  $\epsilon$ )


## neutron detection




## missing mass



## two step



## neutron detection




- ✓ unambiguous detection
- ✓ breakup or resonant decay
- ✓ neutron correlations
- ✗ extremely low efficiency




4 experiments

## missing mass




## two step





## neutron detection




- ✓ unambiguous detection
- ✓ breakup or resonant decay
- ✓ neutron correlations
- ✗ extremely low efficiency




4 experiments

## missing mass




- ✓ detection of 1 charged particle
- ✓ both bound & resonant states
- ✓ mass number well defined
- ✗ insensitive to internal structure
- ✗ cross-section of all protons into  $\textcircled{b}$
- ✗ beam/target contaminant  $\neq \textcircled{a}/\textcircled{c}$




24 experiments

## two step



## neutron detection




- ✓ unambiguous detection
- ✓ breakup or resonant decay
- ✓ neutron correlations
- ✗ extremely low efficiency




4 experiments

## missing mass




- ✓ detection of 1 charged particle
- ✓ both bound & resonant states
- ✓ mass number well defined
- ✗ insensitive to internal structure
- ✗ cross-section of all protons into  $\textcircled{b}$
- ✗ beam/target contaminant  $\neq \textcircled{a}/\textcircled{c}$



24 experiments

## two step




- ✓ detection of 1 charged particle
- ✗ only bound states in second step
- ✗ insensitive to the energy
- ✗ only lower limit of  $A$  inferred
- ✗ contaminant  $\neq \textcircled{a}$  can lead to  $\textcircled{b}$
- ✗ uncontrolled previous step generates huge background, that may lead to  $\textcircled{b}$



8 experiments

## neutron detection



- ✓ unambiguous detection
- ✓ breakup or resonant decay
- ✓ neutron correlations
- ✗ extremely low efficiency



4 experiments


☞ FMM, PRC 65 (2002) 044006

☞ Brill, PL 12 (1964) 51


☞ Koral, NPA 175 (1971) 156

☞ Bystritsky, NIM A834 (2016) 164

## missing mass




- ✓ detection of 1 charged particle
- ✓ both bound & resonant states
- ✓ mass number well defined
- ✗ insensitive to internal structure
- ✗ cross-section of all protons into  $b$
- ✗ beam/target contaminant  $\neq a/c$



24 experiments

☞ Kisamori, PRL 116 (2016) 052501

## two step




- ✓ detection of 1 charged particle
- ✗ only bound states in second step
- ✗ insensitive to the energy
- ✗ only lower limit of  $A$  inferred
- ✗ contaminant  $\neq a$  can lead to  $b$
- ✗ uncontrolled previous step generates huge background, that may lead to  $b$



8 experiments

☞ Détraz, PL 66B (1977) 333



Volume 5, number 4

PHYSICS LETTERS

15 July 1963

## SEARCH FOR A PARTICLE-STABLE TETRA NEUTRON




J. P. SCHIFFER and R. VANDENBOSCH

*Argonne National Laboratory, Argonne, Illinois*

It then seems reasonable that tetra neutrons should be observed inside nuclear reactors in locations where the absorption by nuclei in the moderator is negligible.

As in most experiments of this sort, however, a negative result cannot be regarded as conclusive and further experiments are needed to give additional weight to our result.

We are indebted to Professor R. H. Dalitz for calling this problem to our attention



☞ Schiffer, PL 5 (1963) 292

Volume 5, number 4

PHYSICS LETTERS

15 July 1963

## SEARCH FOR A PARTICLE-STABLE TETRA NEUTRON




J. P. SCHIFFER and R. VANDENBOSCH

*Argonne National Laboratory, Argonne, Illinois*



It then seems reasonable that tetra neutrons should be observed inside nuclear reactors in locations where the absorption by nuclei in the moderator is negligible.

As in most experiments of this sort, however, a negative result cannot be regarded as conclusive and further experiments are needed to give additional weight to our result.

We are indebted to Professor R. H. Dalitz for calling this problem to our attention



☞ Schiffer, PL 5 (1963) 292



☞ Cierjacks, PR 137 (1965) B345

Volume 5, number 4

PHYSICS LETTERS

15 July 1963

## SEARCH FOR A PARTICLE-STABLE TETRA NEUTRON

J. P. SCHIFFER and R. VANDENBOSCH

*Argonne National Laboratory, Argonne, Illinois*

It then seems reasonable that tetra neutrons should be observed inside nuclear reactors in locations where the absorption by nuclei in the moderator is negligible.

As in most experiments of this sort, however, a negative result cannot be regarded as conclusive and further experiments are needed to give additional weight to our result.

We are indebted to Professor R. H. Dalitz for calling this problem to our attention

PHYSICAL REVIEW

VOLUME 137, NUMBER 2B

25 JANUARY 1965





## Further Evidence for the Nonexistence of Particle-Stable Tetraneutrons

S. CIERJACKS, G. MARKUS, W. MICHAELIS, AND W. PÖNITZ

*Institut für Angewandte Kernphysik, Kernforschungszentrum Karlsruhe, Karlsruhe, Germany*

A search for tetraneutrons in the thermal-fission process had a negative result.<sup>8</sup> If tetraneutrons exist at all, the yield in the fast deuteron-induced fission is expected to be about two orders of magnitude higher than in thermal fission. This assumption is reasonable because of the much higher yield of alphas and tritons.<sup>16</sup>

Considering the absence of a Coulomb barrier for the tetraneutron, this particle should occur with a frequency comparable with that of alphas and tritons in spite of the much lower binding energy.<sup>8</sup> Therefore, it seems reasonable to conclude from Table I that the existence of tetraneutrons is most unlikely.



☞ Détraz, PL 66B (1977) 333

Volume 66B, number 4      PHYSICS LETTERS      14 February 1977

## POSSIBLE EXISTENCE OF BOUND NEUTRAL NUCLEI

Claude DÉTRAZ

*Institut de Physique Nucléaire, BP 1, 91406 Orsay, France*

Two neutrons cannot form a bound nuclear system. That does not necessarily imply that several neutrons cannot constitute a bound nucleus. Unfortunately, the neutron-neutron interaction is not known so far with enough precision as to allow a reliable prediction of the binding energy of the lowest state of a multi-neutron system.




None of the experimental searches for bound nuclei of three neutrons [5] or four neutrons [6] were finally successful. Furthermore, the upper limits for the cross sections of the processes in which  ${}^3\text{n}$  or  ${}^4\text{n}$  could have been formed appear small enough to indicate that neither of these nuclei actually exists

This paper reports a search for neutral nuclei heavier than those which were looked for so far. This requests an a-priori abundant source of nuclei such as  ${}^6\text{n}$  or  ${}^8\text{n}$ , and means of detecting them as efficiently as possible.

In view of the apparent failure of more conventional explanations, it is suggested that the observation of  ${}^{72}\text{Zn}$  provides tentative evidence for the existence of bound neutral nuclei

up to mass 9. If  ${}^4\text{n}$  is unbound [6],  ${}^8\text{n}$  and to a lesser degree  ${}^6\text{n}$  appear to be the most likely candidates

# The quest ends at Orsay ?



☞ Détraz, PL 66B (1977) 333

Volume 66B, number 4      PHYSICS LETTERS      14 February 1977

## POSSIBLE EXISTENCE OF BOUND NEUTRAL NUCLEI

Claude DÉTRAZ

*Institut de Physique Nucléaire, BP 1, 91406 Orsay, France*

Two neutrons cannot form a bound nuclear system. That does not necessarily imply that several neutrons cannot constitute a bound nucleus. Unfortunately, the neutron-neutron interaction is not known so far with enough precision as to allow a reliable prediction of the binding energy of the lowest state of a multi-neutron system.

None of the experimental searches for bound nuclei of three neutrons [5] or four neutrons [6] were finally successful. Furthermore, the upper limits for the cross sections of the processes in which  $^3\text{n}$  or  $^4\text{n}$  could have been formed appear small enough to indicate that neither of these nuclei actually exists




This paper reports a search for neutral nuclei heavier than those which were looked for so far. This requests an a-priori abundant source of nuclei such as  $^6\text{n}$  or  $^8\text{n}$ , and means of detecting them as efficiently as possible.

In view of the apparent failure of more conventional explanations, it is suggested that the observation of  $^{72}\text{Zn}$  provides tentative evidence for the existence of bound neutral nuclei

up to mass 9. If  $^4\text{n}$  is unbound [6],  $^8\text{n}$  and to a lesser degree  $^6\text{n}$  appear to be the most likely candidates



# The quest ends at Orsay?



Volume 66B, number 4    PHYSICS LETTERS    14 February 1977

**POSSIBLE EXISTENCE OF BOUND NEUTRAL NUCLEI**

Claude DÉTRAZ  
*Institut de Physique Nucléaire, BP 1, 91406 Orsay, France*

Two neutrons cannot form a bound nuclear system. That does not necessarily imply that several neutrons cannot constitute a bound nucleus. Unfortunately, the neutron-neutron interaction is not known so far with enough precision as to allow a reliable prediction of

VOL 38, NUM 20    PHYSICAL REVIEW LETTERS    16 MAY 1977

**Search for Particle-Bound Polyneutron Systems**

Anthony Turkevich, James R. Cadieux, John Warren, Thanasis Economou, Jerome La Rosa, and H. Roland Heydegger


A search for particle-bound polyneutron systems ( ${}^6n-{}^{12}n$ ) produced in  $\sim 700$ -MeV proton interactions with uranium has yielded negative results. A radiochemical technique was used. The limits on production cross section  $\sim 10^{-3}$  to  $10^{-5} \mu\text{b}$  are in contrast to the positive results reported recently from work with 24-GeV protons on tungsten.

Thus Détraz's polyneutrons either have  $x = 4$ , to which the present experiment is insensitive, or their production has an exceedingly steep energy dependence.

→  ${}^{64-70}\text{Zn}(\underline{A}_n, xn) \underline{{}^{72}\text{Zn}}$  ✓  
 ☞ Détraz, PL 66B (1977) 333

→  $[p+U] \text{}^{208}\text{Pb}(\underline{A}_n, xn) \underline{{}^{212}\text{Pb}}$  ✗  
 ☞ Turkevich, PRL 38 (1977) 1129

al explanations, it is suggested that the observation of  ${}^{72}\text{Zn}$  provides tentative evidence for the existence of bound neutral nuclei up to mass 9. If  ${}^4n$  is unbound [6],  ${}^8n$  and to a lesser degree  ${}^6n$  appear to be the most likely candidates





Volume 66B, number 4    PHYSICS LETTERS    14 February 1977

## POSSIBLE EXISTENCE OF BOUND NEUTRAL NUCLEI

Claude DETRAZ

*Institut de Physique Nucléaire, BP 1, 91406 Orsay, France*

Two neutrons cannot form a bound nuclear system. That does not necessarily imply that several neutrons cannot constitute a bound nucleus. Unfortunately, the neutron-neutron interaction is not known so far with enough precision as to allow a reliable prediction of



VOL 38, NUM 20    PHYSICAL REVIEW LETTERS    16 MAY 1977

## Search for Particle-Bound **Polyneutron** Systems

Anthony Turkevich, James R. Cadieux, John Warren, Thanasis Economou, Jerome La Rosa, and H. Roland Heydegger

A search for particle-bound polyneutron systems ( ${}^6n-{}^{12}n$ ) produced in  $\sim 700$ -MeV proton interactions with uranium has yielded negative results. A radiochemical technique

*Nuclear Physics A350 (1980) 149-156 © North-Holland Publishing Co., Amsterdam*


## THE TETRANEUTRON REVISITED

F.W.N. DE BOER


J.J. VAN RUYVEN, A.W.B. KALSHOVEN and R. VIS

E. SUGARBAKER, C. FIELDS and C.S. ZAIDINS

It seems likely that secondary tritons produced in the (p+W) interactions, with the subsequently induced (t,p) reactions in the detection target, must account for Détraz results. Although shielding against charged fragmentation products had been applied, the number of highly energetic tritons has probably been underestimated<sup>(25)</sup>.




☞ Détraz, PL 66B (1977) 333




☞ Turkevich, PRL 38 (1977) 1129




☞ de Boer, NP A350 (1980) 149



- (4n) Gilly, PL 19 (1965) 335
- (3n) Sperinde, PL 32B (1970) 185
- (3n) Sperinde, NP B78 (1974) 345
- (4n) Ungar, PL 144B (1984) 333 :





(4n) Gilly, PL 19 (1965) 335

(3n) Sperinde, PL 32B (1970) 185

(3n) Sperinde, NP B78 (1974) 345

(4n) Ungar, PL 144B (1984) 333 :


(3n) Jibuti, NP A437 (1985) 687


(3,4n) Stetz, NP A457 (1986) 669

(4n) Goringe, PRC 40 (1989) 2390

(3n) Yuly, PRC 55 (1997) 1848

(3n) Gräter, EPJB 4 (1999) 5 📌





(4n) Gilly, PL 19 (1965) 335

(3n) Sperinde, PL 32B (1970) 185

(3n) Sperinde, NP B78 (1974) 345

(4n) Ungar, PL 144B (1984) 333 :

(3n) Jibuti, NP A437 (1985) 687



(3,4n) Stetz, NP A457 (1986) 669


(4n) Goringe, PRC 40 (1989) 2390

(3n) Yuly, PRC 55 (1997) 1848

(3n) Gräter, EPJB 4 (1999) 5

(4n) Chultem, NP A316 (1979) 290 :





(4n) Gilly, PL 19 (1965) 335

(3n) Sperinde, PL 32B (1970) 185

(3n) Sperinde, NP B78 (1974) 345

(4n) Ungar, PL 144B (1984) 333 :

(3n) Jibuti, NP A437 (1985) 687


(3,4n) Stetz, NP A457 (1986) 669

(4n) Goringe, PRC 40 (1989) 2390



(3n) Yuly, PRC 55 (1997) 1848

(3n) Gräter, EPJB 4 (1999) 5

(4n) Chultem, NP A316 (1979) 290 :



(3n) Gornov, NP A531 (1991) 613 :



→ “phase-space can lead to a **distortion** of the results”  
 → “the rather **poor** experimental data” ...

## Photon Spectrum in Pion Capture on Tritium<sup>†</sup>

J. A. Bistirlich, S. Cooper, K. M. Crowe, and F. T. Shively<sup>†</sup>

*Lawrence Berkeley Laboratory, University of California, Berkeley, California 94720*

E. R. Grilly, J. P. Perroud,<sup>‡</sup> and R. H. Sherman

*Los Alamos Scientific Laboratory, Los Alamos, New Mexico 87545*

H. W. Baer<sup>§</sup>

*Case Western Reserve University, Cleveland, Ohio 44106*

P. Truöl

*Physik-Institut der Universität, Zürich, Switzerland*




FIG. 1. The experimental setup at LAMPF showing the pair spectrometer and liquid-tritium target. The inset shows a cross section of the target cell obtained from an x-ray radiograph.

The overall fit to the data is satisfactory, although small excesses of events in the low-mass region  $7 < E_x(3n) \leq 16$  MeV are observed. Considering the low statistics and uncertainty in background subtraction, it would be premature to regard this as evidence for a  $T = \frac{3}{2}$  resonance




FIG. 2. (a) Raw photon spectrum obtained from the tritium target. The inset shows our resolution obtained at 129.4 MeV. (b) Background spectra for hydrogen, deuterium, steel, and CH. (c) Spectrum from reaction  $\pi^- + {}^3\text{H} \rightarrow n + n + n + \gamma$  after subtraction of  ${}^1\text{H}$ ,  ${}^2\text{H}$ , steel, and scintillator contributions. Solid curve is the theoretical spectrum of Phillips and Roig (Ref. 10) (see text), folded with acceptance and instrumental line shape and normalized to the total number of photons.

## Photon Spectrum in Pion Capture on Tritium†

J. A. Bistirlich, S. Cooper, K. M. Crowe, and F. T. Shively†

*Lawrence Berkeley Laboratory, University of California, Berkeley, California 94720*

E. R. Grilly, J. P. Perroud,‡ and R. H. Sherman

*Los Alamos Scientific Laboratory, Los Alamos, New Mexico 87545*

H. W. Baer§

*Case Western Reserve University, Cleveland, Ohio 44106*

P. Truöl

*Physik-Institut der Universität, Zürich, Switzerland*




FIG. 1. The experimental setup at LAMPF showing the pair spectrometer and liquid-tritium target. The inset shows a cross section of the target cell obtained from an x-ray radiograph.

The overall fit to the data is satisfactory, although small excesses of events in the low-mass region  $7 < E_x(3n) \leq 16$  MeV are observed. Considering the low statistics and uncertainty in background subtraction, it would be premature to regard this as evidence for a  $T = \frac{3}{2}$  resonance




FIG. 2. (a) Raw photon spectrum obtained from the tritium target. The inset shows our resolution obtained at 129.4 MeV. (b) Background spectra for hydrogen, deuterium, steel, and CH. (c) Spectrum from reaction  $\pi^- + {}^3\text{H} \rightarrow n + n + n + \gamma$  after subtraction of  ${}^1\text{H}$ ,  ${}^2\text{H}$ , steel, and scintillator contributions. Solid curve is the theoretical spectrum of Phillips and Roig (Ref. 10) (see text), folded with acceptance and instrumental line shape and normalized to the total number of photons.



## Photon Spectrum in Pion Capture on Tritium†

J. A. Bistirlich, S. Cooper, K. M. Crowe, and F. T. Shively†  
Lawrence Berkeley Laboratory, University of California, Berkeley, California 94720

E. R. Grilly, J. P. Perroud,‡ and R. H. Sherman  
Los Alamos Scientific Laboratory, Los Alamos, New Mexico 87545

H. W. Baer§  
Case Western Reserve University, Cleveland, Ohio 44106

P. Truöl  
Physik-Institut der Universität, Zürich, Switzerland




FIG. 1. The experimental setup at LAMPF showing the pair spectrometer and liquid-tritium target. The inset shows a cross section of the target cell obtained from an x-ray radiograph.

The overall fit to the data is satisfactory, although small excesses of events in the low-mass region  $7 < E_x(3n) \leq 16$  MeV are observed. Considering the low statistics and uncertainty in background subtraction, it would be premature to regard this as evidence for a  $T = \frac{3}{2}$  resonance




FIG. 2. (a) Raw photon spectrum obtained at 129.4 MeV. (b) Background spectra for hydrogen, deuterium, steel, and CH. (c) Spectrum after subtraction of background for tritium, showing 1064 events. The red circle highlights a peak at 129.4 MeV. The x-axis is photon energy  $E_\gamma$  (MeV) from 50 to 140.

## UPPER LIMITS FOR BOUND STATES AND RESONANCE BEHAVIOR IN THE TRINEUTRON SYSTEM

J. P. MILLER †, J. A. BISTIRLICH, K. M. CROWE, S. S. ROSENBLUM,  
P. C. ROWE and F. T. SHIVELY

Lawrence Berkeley Laboratory, University of California, Berkeley, California 94720, USA

E. R. GRILLY, E. C. KERR, J. NOVAK and R. H. SHERMAN  
Los Alamos Scientific Laboratory, Los Alamos, New Mexico 87545, USA

H. BRÄNDLE \*\*  
Physics Department, University of California, Los Angeles, California 90024, USA

G. STRASSNER \*\*\* and P. TRUÖL  
Physik-Institut der Universität Zürich, CH-8001 Zürich, Switzerland




Fig. 4. Measured tritium spectrum with background subtracted. Curve a, Amado model and curve b, plane wave for 3n final state, from refs. 3,21)

In conclusion, we have performed an experiment expected to be highly sensitive to the presence of 3n structure near threshold and see no evidence for it, other than a very pronounced shift to low 3n energy which can be explained in terms of the simple s-wave pairwise interaction between neutrons in the final state.

## Photon Spectrum in Pion Capture on Tritium†

J. A. Bistirlich, S. Cooper, K. M. Crowe, and F. T. Shively†  
Lawrence Berkeley Laboratory, University of California, Berkeley, California 94720

E. R. Grilly, J. P. Perroud,‡ and R. H. Sherman  
Los Alamos Scientific Laboratory, Los Alamos, New Mexico 87545

H. W. Baer§  
Case Western Reserve University, Cleveland, Ohio 44106

P. Truöl  
Physik-Institut der Universität, Zürich, Switzerland




FIG. 1. The experimental setup at LAMPF showing the pair spectrometer and liquid-tritium target. The inset shows a cross section of the target cell obtained from an x-ray radiograph.

The overall fit to the data is satisfactory, although small excesses of events in the low-mass region  $7 < E_x(3n) \leq 16$  MeV are observed. Considering the low statistics and uncertainty in background subtraction, it would be premature to regard this as evidence for a  $T = \frac{3}{2}$  resonance




FIG. 2. (a) Raw photon spectrum obtained at tritium target. The inset shows our result retained at 129.4 MeV. (b) Background spectra for hydrogen, deuterium, steel, and CH. (c) Spectrum after subtraction of backgrounds for tritium, steel, and scintillator contributions. Theoretical spectrum of Phillips et al. (see text), folded with acceptance and shape and normalized to the total number of events.

## UPPER LIMITS FOR BOUND STATES AND RESONANCE BEHAVIOR IN THE TRINEUTRON SYSTEM

J. P. MILLER †, J. A. BISTIRLICH, K. M. CROWE, S. S. ROSENBLUM,  
P. C. ROWE and F. T. SHIVELY

Lawrence Berkeley Laboratory, University of California, Berkeley, California 94720, USA

E. R. GRILLY, E. C. KERR, J. NOVAK and R. H. SHERMAN  
Los Alamos Scientific Laboratory, Los Alamos, New Mexico 87545, USA

H. BRÄNDLE ††  
Physics Department, University of California, Los Angeles, California 90024, USA

G. STRASSNER ††† and P. TRUÖL  
Physik-Institut der Universität Zürich, CH-8001 Zürich, Switzerland




Fig. 4. Measured tritium spectrum with background subtracted. Curve a, Amado model and curve b, plane wave for 3n final state, from refs. 3,21)

In conclusion, we have performed an experiment expected to be highly sensitive to the presence of 3n structure near threshold and see no evidence for it, other than a very pronounced shift to low 3n energy which can be explained in terms of the simple s-wave pairwise interaction between neutrons in the final state.

☞ Ajdačić, PRL 14 (1965) 444 :  ${}^3\text{H}(n, p){}^3\text{n}$  (✓)

☞ Thornton, PRL 17 (1966) 701 :  ${}^3\text{H}(n, p){}^3\text{n}$  ✗

☞ Ohlsen, PR 176 (1968) 1163 :  ${}^3\text{H}(t, {}^3\text{He})3\text{n}$  (✓)

→ very poor data

→ some unclear “enhancements” ...

☞ Ajdačić, PRL 14 (1965) 444 :  ${}^3\text{H}(n, p){}_3\text{n}$  (✓)

☞ Thornton, PRL 17 (1966) 701 :  ${}^3\text{H}(n, p){}_3\text{n}$  ✗

☞ Ohlsen, PR 176 (1968) 1163 :  ${}^3\text{H}(t, {}^3\text{He}){}_3\text{n}$  (✓)


→ very poor data

→ some unclear “enhancements” ...

☞ Cerny, PL 53B (1974) 247 :

${}^7\text{Li}({}^7\text{Li}, {}^{11}\text{C}){}_3\text{n}$  ✗

${}^7\text{Li}({}^7\text{Li}, {}^{10}\text{C}){}_4\text{n}$  ✗



Ajdačić, PRL 14 (1965) 444 :  ${}^3\text{H}(n, p){}_3\text{n}$  (✓)

Thornton, PRL 17 (1966) 701 :  ${}^3\text{H}(n, p){}_3\text{n}$  ✗

Ohlsen, PR 176 (1968) 1163 :  ${}^3\text{H}(t, {}^3\text{He}){}_3\text{n}$  (✓)

→ very poor data

→ some unclear “enhancements” ...

Belozyorov, NP A477 (1988) 131 :

${}^7\text{Li}({}^{11}\text{B}, {}^{15}\text{O}){}_3\text{n}$  ✗

${}^7\text{Li}({}^9\text{Be}, {}^{12}\text{N}){}_4\text{n}$  ✗


${}^7\text{Li}({}^{11}\text{B}, {}^{14}\text{O}){}_4\text{n}$  ✗

${}^9\text{Be}({}^9\text{Be}, {}^{14}\text{O}){}_4\text{n}$  ✗

Cerny, PL 53B (1974) 247 :

${}^7\text{Li}({}^7\text{Li}, {}^{11}\text{C}){}_3\text{n}$  ✗

${}^7\text{Li}({}^7\text{Li}, {}^{10}\text{C}){}_4\text{n}$  ✗



# Transfer : exploring beam/target combinations

Ajdačić, PRL 14 (1965) 444 :  ${}^3\text{H}(n, p){}_3\text{n}$  (✓)

Thornton, PRL 17 (1966) 701 :  ${}^3\text{H}(n, p){}_3\text{n}$  ✗

Ohlsen, PR 176 (1968) 1163 :  ${}^3\text{H}(t, {}^3\text{He}){}_3\text{n}$  (✓)


→ very poor data

→ some unclear “enhancements” ...

Cerny, PL 53B (1974) 247 :

${}^7\text{Li}({}^7\text{Li}, {}^{11}\text{C}){}_3\text{n}$  ✗

${}^7\text{Li}({}^7\text{Li}, {}^{10}\text{C}){}_4\text{n}$  ✗





Belozyorov, NP A477 (1988) 131 :

${}^7\text{Li}({}^{11}\text{B}, {}^{15}\text{O}){}_3\text{n}$  ✗       ${}^7\text{Li}({}^9\text{Be}, {}^{12}\text{N}){}_4\text{n}$  ✗

${}^7\text{Li}({}^{11}\text{B}, {}^{14}\text{O}){}_4\text{n}$  ✗       ${}^9\text{Be}({}^9\text{Be}, {}^{14}\text{O}){}_4\text{n}$  ✗

Bohlen, NP A583 (1995) 775 :



# Transfer : exploring beam/target combinations

Ajdačić, PRL 14 (1965) 444 :  ${}^3\text{H}(n, p){}_3\text{n}$  (✓)

Thornton, PRL 17 (1966) 701 :  ${}^3\text{H}(n, p){}_3\text{n}$  ✗

Ohlsen, PR 176 (1968) 1163 :  ${}^3\text{H}(t, {}^3\text{He}){}_3\text{n}$  (✓)


→ very poor data

→ some unclear “enhancements” ...

Cerny, PL 53B (1974) 247 :

${}^7\text{Li}({}^7\text{Li}, {}^{11}\text{C}){}_3\text{n}$  ✗

${}^7\text{Li}({}^7\text{Li}, {}^{10}\text{C}){}_4\text{n}$  ✗




Belozyorov, NP A477 (1988) 131 :

${}^7\text{Li}({}^{11}\text{B}, {}^{15}\text{O}){}_3\text{n}$  ✗       ${}^7\text{Li}({}^9\text{Be}, {}^{12}\text{N}){}_4\text{n}$  ✗

${}^7\text{Li}({}^{11}\text{B}, {}^{14}\text{O}){}_4\text{n}$  ✗       ${}^9\text{Be}({}^9\text{Be}, {}^{14}\text{O}){}_4\text{n}$  ✗



Bohlen, NP A583 (1995) 775 :

$[{}^6\text{Li}_2\text{O}/\text{CD}_2]({}^{14}\text{C}, {}^{13}\text{N}) \equiv -\text{red circle} + \text{blue circles}$



Aleksandrov, JETPL 81-2 (2005) 43 : confirms Cerny's work


# Sculpting exotic beams (SAMURAI21)




$\beta \sim 0.6c$  !!!














# Sculpting exotic beams (SAMURAI21)



# Sculpting exotic beams (SAMURAI21)







► Kisamori, PRL 116 (2016) 052501 →  $E(4n) = 0.8 \pm 1.3$  MeV

► FMM, PRC 65 (2002) 044006  
 ► FMM, arXiv nucl-ex/0504009 }  $E(4n) \in [-1, +2]$  MeV


► ‘Exact’ calculations are categorical!

☞ Glöckle, PRC 18 (1978) 564 :  $V_{nn} \times 4.2$

☞ Offermann, NPA 318 (1979) 138 :  $V_{nn} \times 3.7$  (+P-waves)

☞ Witała, PRC 60 (1999) 024002 : avoid  ${}^2n$  with  $V_{nn}({}^1S_0) \times 1$


☞ Hemmdan, PRC 66 (2002) 054001 :



“ ${}^3n$  resonances close to the physical region will not exist”

► ‘Exact’ calculations are categorical!

- ☞ Glöckle, PRC 18 (1978) 564 :  $V_{nn} \times 4.2$
- ☞ Offermann, NPA 318 (1979) 138 :  $V_{nn} \times 3.7$  (+P-waves)
- ☞ Witała, PRC 60 (1999) 024002 : avoid  ${}^2n$  with  $V_{nn}({}^1S_0) \times 1$
- ☞ Hemmdan, PRC 66 (2002) 054001 :




“ $3n$  resonances close to the physical region will not exist”

- (3n) ☞ Lazauskas, PRC 71 (2005) 044004 : 3NF ✗
- (4n) ☞ Lazauskas, PRC 72 (2005) 034003 : 4NF ✗
- (3,4n) ☞ Hiyama, PRC 93 (2016) 044004 : 3NF( $T=3/2$ ) ✗!

# Theory: another hard & interesting quest

► ‘Exact’ calculations are categorical!

- ☞ Glöckle, PRC 18 (1978) 564 :  $V_{nn} \times 4.2$
- ☞ Offermann, NPA 318 (1979) 138 :  $V_{nn} \times 3.7$  (+P-waves)
- ☞ Witała, PRC 60 (1999) 024002 : avoid  ${}^2n$  with  $V_{nn}({}^1S_0) \times 1$
- ☞ Hemmdan, PRC 66 (2002) 054001 :




“ $3n$  resonances close to the physical region will not exist”

- (3n) ☞ Lazauskas, PRC 71 (2005) 044004 : 3NF ✗
- (4n) ☞ Lazauskas, PRC 72 (2005) 034003 : 4NF ✗
- (3,4n) ☞ Hiyama, PRC 93 (2016) 044004 : 3NF( $T=3/2$ ) ✗!

► Many-body approximations, not so much ...

- ☞ Pieper, PRL 90 (2003), 252501 :




“the resonance, if it exists at all, must be very broad”

- ☞ Shirokov, PRL 117 (2016) 182502
  - ☞ Gandolfi, PRL 118 (2017) 232501
  - ☞ Fosse, PRL 119 (2017) 032501
  - ☞ Li, PRC 100 (2019) 054313
- }  $3n/4n$  ✓?

# Theory: another hard & interesting quest

► ‘Exact’ calculations are categorical!

- Glöckle, PRC 18 (1978) 564 :  $V_{nn} \times 4.2$
- Offermann, NPA 318 (1979) 138 :  $V_{nn} \times 3.7$  (+P-waves)
- Witała, PRC 60 (1999) 024002 : avoid  ${}^2n$  with  $V_{nn}({}^1S_0) \times 1$
- Hemmdan, PRC 66 (2002) 054001 :




“ ${}^3n$  resonances close to the physical region will not exist”

- (3n) Lazauskas, PRC 71 (2005) 044004 : 3NF ✗
- (4n) Lazauskas, PRC 72 (2005) 034003 : 4NF ✗
- (3,4n) Hiyama, PRC 93 (2016) 044004 : 3NF( $T=3/2$ ) ✗!

► Many-body approximations, not so much ...

Pieper, PRL 90 (2003), 252501 :




“the resonance, if it exists at all, must be very broad”

- Shirokov, PRL 117 (2016) 182502
  - Gandolfi, PRL 118 (2017) 232501
  - Fosse, PRL 119 (2017) 032501
  - Li, PRC 100 (2019) 054313
- }  ${}^3n/{}^4n$  ✓?

- Deltuva, PRL 123 (2019) 069201
  - Deltuva, PRC 100 (2019) 044002
  - Ishikawa, PRC 102 (2020) 034002
- }  ${}^3n/{}^4n$  ✗ !!!  
(trap/evolution/scaling)
- Deltuva, PLB 782 (2018) 238
  - Higgins, PRL 125 (2020) 052501
- } QM enhancements ...






- ▶ 36 works published !
  - 14 exclusively for tetraneutron
  - 3 positive signals !
    - 1 strong but refuted
    - 2 weak but uncontested (yet)




☞ Détraz, PL 66B (1977) 333



☞ FMM, PRC 65 (2002) 044006



☞ Kisamori, PRL 116 (2016) 052501




- ▶ 36 works published !
  - 14 exclusively for **tetraneutron**
  - 3 positive signals !
    - 1 strong but refuted
    - **2 weak** but uncontested (yet)
  - recurring pattern in XX century :
    - **qualitative** increase in 2020s ?



☞ Détraz, PL 66B (1977) 333



☞ FMM, PRC 65 (2002) 044006



☞ Kisamori, PRL 116 (2016) 052501

- ▶ Three experiments: same beam ( $^8\text{He}$ ) & energy (150–200 MeV/N)?

reaction	initial state	final state	$\sigma$	results
('16) $^4\text{He} (^8\text{He}, \alpha\alpha) ^4\text{n}$ Shimoura, NP1512-SHARAQ10			nb	$N_{\text{evt}} \sim 10\text{ s}$ $^4\text{n}: E, \Gamma$
('17) $^8\text{He} (p, p\alpha) ^4\text{n}$ Paschalis, NP1406-SAMURAI19			$\mu\text{b}$	$N_{\text{evt}} \sim 1000\text{ s}$ $^4\text{n}: E, \Gamma$
('17) $^8\text{He} (p, 2p) \{^3\text{H} + ^4\text{n}\}$ FMM/Yang, NP1512-SAMURAI34			mb	$N_{\text{evt}} \sim 10,000\text{ s}$ $^4\text{n} \& ^7\text{H}: E, \Gamma, \Omega$

► Three experiments: same beam ( $^8\text{He}$ ) & energy (150–200 MeV/N)?

reaction	initial state	final state	$\sigma$	results
('16) $^4\text{He} (^8\text{He}, \alpha\alpha) ^4\text{n}$ Shimoura, NP1512-SHARAQ10			nb	$N_{\text{evt}} \sim 10\text{ s}$ $^4\text{n}: E, \Gamma$
('17) $^8\text{He} (p, p\alpha) ^4\text{n}$ Paschalis, NP1406-SAMURAI19			$\mu\text{b}$	$N_{\text{evt}} \sim 1000\text{ s}$ $^4\text{n}: E, \Gamma$
('17) $^8\text{He} (p, 2p) \{^3\text{H} + ^4\text{n}\}$ FMM/Yang, NP1512-SAMURAI34			mb	$N_{\text{evt}} \sim 10,000\text{ s}$ $^4\text{n} \& ^7\text{H}: E, \Gamma, \Omega$

→ very 'simple' formula:

$$N_{\text{evt}} \propto I \times \sigma \times \rho \times \epsilon$$

{


beam **intensity**

reaction **cross-section**


number of **target nuclei**

detection **efficiency**

# Hydrogen 7 & Tetraneutron 'emission' ?




►  ${}^8\text{He} (p, 2p) {}^7\text{H}$  @ 150 MeV/N:




→ 7-body final state!


# Hydrogen 7 & Tetraneutron 'emission' ?



- $N = 6$  ( $p_{3/2}$ ) sub-shell closure ?




►  ${}^8\text{He}(p,2p){}^7\text{H}$  @ 150 MeV/N:




→ 7-body final state!

# Hydrogen 7 & Tetra-neutron 'emission' ?




- $N = 6$  ( $p_{3/2}$ ) sub-shell closure ?



- an array of arrays :

- MINOS liquid H target
- DALI NaI crystals


▶  ${}^8\text{He}(p,2p){}^7\text{H}$  @ 150 MeV/N :




→ 7-body final state !

→ FWHM ~ few MeV

# Hydrogen 7 & Tetra-neutron 'emission' ?




- $N = 6$  ( $p_{3/2}$ ) sub-shell closure ?



- an array of arrays :

- MINOS liquid H target
- DALI NaI crystals
- SAMURAI spectrometer
- NEBULA + NeuLAND

►  ${}^8\text{He}(p,2p){}^7\text{H}$  @ 150 MeV/N :




→ 7-body final state !


→ FWHM ~ few MeV → 100 keV !



# Hydrogen 7 & Tetraneutron 'emission' ?




- N = 6 (p<sub>3/2</sub>) sub-shell closure ?



- an array of arrays :


- MINOS liquid H target
- DALI NaI crystals
- SAMURAI spectrometer
- NEBULA + NeuLAND


► <sup>8</sup>He (p,2p) <sup>7</sup>H @ 150 MeV/N :






- 7-body final state !
- FWHM ~ few MeV → 100 keV !
- (2p+t+3n) ~ 150 keV

- angular correlations → E(4n) !
















NeuLAND

NEBULA




NeuLAND

NEBULA




☞ Schiffer (1963) “As in most experiments of this sort, however, a negative result cannot be regarded as conclusive”



► Extraordinary series of experiments !

- fascinating ideas
- some precise  $3n$  results
- few weak  $4n$  signals

📖 Schiffer (1963) *“As in most experiments of this sort, however, a negative result cannot be regarded as conclusive”*




► Extraordinary series of experiments !

- fascinating ideas
- some precise  $3n$  results
- few weak  $4n$  signals :
  - refute / states / “enhancements” ?
- promising 2020s perspectives !
  - high statistics & resolution
  - first  $6n$  experiments ( $^{10}\text{He}$  decay) ...



☞ Schiffer (1963) “As in most experiments of this sort, however, a negative result cannot be regarded as conclusive”




► Extraordinary series of experiments !

- fascinating ideas
- some precise  $3n$  results
- few weak  $4n$  signals :  
→ refute / states / “enhancements” ?
- promising 2020s perspectives !  
→ high statistics & resolution  
→ first  $6n$  experiments ( $^{10}\text{He}$  decay) ...


► Some theoretical certainties :

- all exact calculations categorical :  
→ signals cannot be  $3/4n$  “states” !



- many-body approx. claim  $3/4n$  resonances ...
- consensus : independently of  $V_{nn(n)}$  !  
→ trap & global scaling : thresholds  
→ extrapolation of states into continuum

☞ Schiffer (1963) “As in most experiments of this sort, however, a negative result cannot be regarded as conclusive”




▶ Extraordinary series of experiments !

- fascinating ideas
- some precise  $3n$  results
- few weak  $4n$  signals :  
→ refute / states / “enhancements” ?
- promising 2020s perspectives !  
→ high statistics & resolution  
→ first  $6n$  experiments ( $^{10}\text{He}$  decay) ...

▶ Some theoretical certainties :

- all exact calculations categorical :  
→ signals cannot be  $3/4n$  “states” !



- many-body approx. claim  $3/4n$  resonances ...
- consensus : independently of  $V_{nn(n)}$  !  
→ trap & global scaling : thresholds  
→ extrapolation of states into continuum

▶ Some theoretical hopes ?

- benchmark experimental results !
- QM “enhancements” ?  $6,8,10n$  trends ?

## Reaction Seminars 2021 (April 1, 2021)









**From one to many : a neutral history**

F. Miguel Marqués




► Expand NEBULA multineutron capabilities :


- France: LPC, IRFU, IPNO
- Japan: TITech, RIKEN
- +90 bars: Commissioning & Day-1 in 2021 (?)




$E_d$  [MeV] ( $^{24+x}\text{O} \rightarrow ^{24}\text{O} + xn$ )



Kurchatov Institute



## Kurchatov Institute



ISSN 0021-3640, JETP Letters, 2012, Vol. 96, No. 5, pp. 280–284. © Pleiades Publishing, Inc., 2012.

### Possible Observation of Light Neutron Nuclei in the Alpha-Particle-Induced Fission of $^{238}\text{U}$

B. G. Novatsky, E. Yu. Nikolsky, S. B. Sakuta, and D. N. Stepanov

National Research Centre Kurchatov Institute, pl. Akademika Kurchatova 1, Moscow, 123182 Russia





Fig. 1. (a) Measured gamma-ray spectrum of a  $^{88}\text{SrCO}_3$  sample irradiated with products of  $^{238}\text{U}$  fission induced by alpha particles (the most intense lines are shown—see main body of the text). (b) Segment of this gamma-ray spectrum in the energy range of 1300–1500 keV. The arrow indicates the  $^{92}\text{Sr}$  (1384 keV) gamma line.

The formation of this nucleus was associated with a four-neutron-transferring reaction involving a nuclear-stable multineutron:  $^{88}\text{Sr}(^x n, (x-4)n)^{92}\text{Sr}$ . In order to confirm this result, it is necessary to perform further experiments with heavier bombarding particles ( $^{11}\text{B}$  and  $^{12}\text{C}$ ) and with other activated targets.



## Kurchatov Institute



ISSN 0021-3640, JETP Letters, 2012, Vol. 96, No. 5, pp. 280–284. © Pleiades Publishing, Inc., 2012.

### Possible Observation of Light Neutron Nuclei in the Alpha-Particle-Induced Fission of $^{238}\text{U}$

B. G. Novatsky, E. Yu. Nikolsky, S. B. Sakuta, and D. N. Stepanov

National Research Centre Kurchatov Institute, pl. Akademika Kurchatova 1, Moscow, 123182 Russia





Fig. 1. (a) Measured gamma-ray spectrum of a  $^{88}\text{SrCO}_3$  sample irradiated with products of  $^{238}\text{U}$  fission induced by alpha particles (the most intense lines are shown—see main body of the text). (b) Segment of this gamma-ray spectrum in the energy range of 1300–1500 keV. The arrow indicates the  $^{92}\text{Sr}$  (1384 keV) gamma line.

The formation of this nucleus was associated with a four-neutron-transferring reaction involving a nuclear-stable multineutron:  $^{88}\text{Sr}(^x n, (x-4)n)^{92}\text{Sr}$ .

In order to confirm this result, it is necessary to perform further experiments with heavier bombarding particles ( $^{11}\text{B}$  and  $^{12}\text{C}$ ) and with other activated targets.



## Kurchatov Institute



ISSN 0021-3640, JETP Letters, 2012, Vol. 96, No. 5, pp. 280–284. © Pleiades Publishing, Inc., 2012.

### Possible Observation of Light Neutron Nuclei in the Alpha-Particle-Induced Fission of $^{238}\text{U}$

B. G. Novatsky, E. Yu. Nikolsky, S. B. Sakuta, and D. N. Stepanov

National Research Centre Kurchatov Institute, pl. Akademika Kurchatova 1, Moscow, 123182 Russia




Fig. 1. (a) Measured gamma-ray spectrum of a  $^{88}\text{SrCO}_3$  sample irradiated with products of  $^{238}\text{U}$  fission induced by alpha particles (the most intense lines are shown—see main body of the text). (b) Segment of this gamma-ray spectrum in the energy range of 1300–1500 keV. The arrow indicates the  $^{92}\text{Sr}$  (1384 keV) gamma line.

The formation of this nucleus was associated with a four-neutron-transfer reaction involving a nuclear-stable multineutron:  $^{88}\text{Sr}(^x n, (x-4)n)^{92}\text{Sr}$ .

In order to confirm this result, it is necessary to perform further experiments with heavier bombarding particles ( $^{11}\text{B}$  and  $^{12}\text{C}$ ) and with other activated targets.



### Detection of Light Neutron Nuclei in the Alpha-Particle-Induced Fission of $^{238}\text{U}$ by the Activation Method with $^{27}\text{Al}$

B. G. Novatsky, S. B. Sakuta\*, and D. N. Stepanov

National Research Centre Kurchatov Institute, pl. Akademika Kurchatova 1, Moscow, 123182 Russia

ISSN 0021-3640, JETP Letters, 2013, Vol. 98, No. 11, pp. 656–660. © Pleiades Publishing, Inc., 2013.




Fig. 2. (Color online) (a) Energy spectrum of gamma rays from the  $^{27}\text{Al}$  sample that was irradiated by the products of alpha-particle-induced fission of the  $^{238}\text{U}$  nucleus. (b) Fragment of this gamma-ray spectrum in the energy range of 1330–2250 keV. The arrows mark the 1342- and 1779-keV gamma lines from the beta decay of  $^{28}\text{Mg}$  and  $^{28}\text{Al}$  nuclei, respectively.

The results of two independent experiments indicate that nuclear-stable multineutrons (most likely,  $^6n$ ) are emitted from the alpha-particle-induced ternary fission of  $^{238}\text{U}$ . In the future, we are going to improve the statistics of the measurements by increasing the intensity of the beam and irradiation time of sample.




Novatsky, JETPL 98-11 (2013) 656

Novatsky, JETPL 96-5 (2012) 280



## Kurchatov Institute



ISSN 0021-3640, JETP Letters, 2012, Vol. 96, No. 5, pp. 280–284. © Pleiades Publishing, Inc., 2012.

### Possible Observation of Light Neutron Nuclei in the Alpha-Particle-Induced Fission of $^{238}\text{U}$

B. G. Novatsky, E. Yu. Nikolsky, S. B. Sakuta, and D. N. Stepanov

National Research Centre Kurchatov Institute, pl. Akademika Kurchatova 1, Moscow, 123182 Russia




Fig. 1. (a) Measured gamma-ray spectrum of a  $^{88}\text{SrCO}_3$  sample irradiated with products of  $^{238}\text{U}$  fission induced by alpha particles (the most intense lines are shown—see main body of the text). (b) Segment of this gamma-ray spectrum in the energy range of 1300–1500 keV. The arrow indicates the  $^{92}\text{Sr}$  (1384 keV) gamma line.

The formation of this nucleus was associated with a four-neutron-transfer reaction involving a nuclear-stable multineutron:  $^{88}\text{Sr}(^x n, (x-4)n)^{92}\text{Sr}$ .

In order to confirm this result, it is necessary to perform further experiments with heavier bombarding particles ( $^{11}\text{B}$  and  $^{12}\text{C}$ ) and with other activated targets.



### Detection of Light Neutron Nuclei in the Alpha-Particle-Induced Fission of $^{238}\text{U}$ by the Activation Method with $^{27}\text{Al}$

B. G. Novatsky, S. B. Sakuta\*, and D. N. Stepanov

National Research Centre Kurchatov Institute, pl. Akademika Kurchatova 1, Moscow, 123182 Russia

ISSN 0021-3640, JETP Letters, 2013, Vol. 98, No. 11, pp. 656–660. © Pleiades Publishing, Inc., 2013.




Fig. 2. (Color online) (a) Energy spectrum of gamma rays from the  $^{27}\text{Al}$  sample that was irradiated by the products of alpha-particle-induced fission of the  $^{238}\text{U}$  nucleus. (b) Fragment of this gamma-ray spectrum in the energy range of 1330–2250 keV. The arrows mark the 1342- and 1779-keV gamma lines from the beta decay of  $^{28}\text{Mg}$  and  $^{28}\text{Al}$  nuclei, respectively.

The results of two independent experiments indicate that nuclear-stable multineutrons (most likely,  $^6n$ ) are emitted from the alpha-particle-induced ternary fission of  $^{238}\text{U}$ . In the future, we are going to improve the statistics of the measurements by increasing the intensity of the beam and irradiation time of sample.



Novatsky, JETPL 98-11 (2013) 656

Novatsky, JETPL 96-5 (2012) 280